

A SEMI-FINITE ALGEBRA ASSOCIATED TO A SUBFACTOR PLANAR ALGEBRA.

A. GUIONNET*, V. JONES† AND D. SHLYAKHTENKO‡

ABSTRACT. We canonically associate to any planar algebra two type II_∞ factors $\mathfrak{M}_\pm$. The subfactors constructed previously by the authors in [GJS08] are isomorphic to compressions of $\mathfrak{M}_\pm$ to finite projections. We show that each $\mathfrak{M}_\pm$ is isomorphic to an amalgamated free product of type I von Neumann algebras with amalgamation over a fixed discrete type I von Neumann subalgebra. In the finite-depth case, existing results in the literature imply that $\mathfrak{M}_+ \cong \mathfrak{M}_-$ is the amplification a free group factor on a finite number of generators. As an application, we show that the factors M_j constructed in [GJS08] are isomorphic to interpolated free group factors $L(\mathbb{F}(r_j))$, $r_j = 1 + 2\delta^{-2j}(\delta - 1)I$, where δ^2 is the index of the planar algebra and I is its global index. Other applications include computations of laws of Jones-Wenzl projections.

1. INTRODUCTION.

In this paper, we associate a pair of semi-finite von Neumann algebras $\mathfrak{M}_\pm$ to a planar algebra $\mathcal{P}$. The algebras $\mathfrak{M}_\pm$ are obtained via the GNS construction from a certain non-unital tracial inductive limit algebra V_+ which arises canonically from $\mathcal{P}$. These algebras have an interesting structure, and the paper is mainly devoted to their study.

To state our main application, let $\mathcal{P}$ be a subfactor planar algebra of index δ^2 , and let us denote by $M_k = M_k(\mathcal{P})$ the von Neumann algebra generated in the GNS representation of $(\mathcal{P}, \wedge_k, Tr_k)$ (see Def. 7 and 8 in [GJS08]). We prove:

Theorem 1. *Assume that $\mathcal{P}$ is finite-depth with global index I . Then $M_k \cong L(\mathbb{F}(r_k))$ with $r_k = 1 + 2\delta^{-2k}(\delta - 1)I$.*

We refer the reader to [Ocn88, EK98] for the definition of global index I . If Γ is the principal graph of $\mathcal{P}$ and μ is the Perron-Frobenius eigenvector normalized by $\mu(*) = 1$, then $I = \frac{1}{2} \sum_{v \in \Gamma} \mu(v)^2$. This formula is consistent with the result of Kodiyalam and Sunder in the depth two case [KSa].

The main step in proving 1 is to prove that the amplifications of M_k are isomorphic to type II_∞ von Neumann algebra $\mathfrak{M}_+$ or $\mathfrak{M}_-$ (the choice of sign is according to the parity of k). It turns out that each $\mathfrak{M}_\pm$ admits a description as a (possibly infinite) free product with amalgamation over a discrete type I von Neumann subalgebra of type I von Neumann algebras. In the finite-depth case, this is sufficient to determine the isomorphism class of $\mathfrak{M}_\pm$ using the work of Dykema [BDJ06, Dyk93, Dyk94, Dyk09].

We note that r_k the statement of Theorem 1 satisfy $(r_k - 1) = \delta^2(r_{k+1} - 1)$.

*UMPA, ENS Lyon, 46 allée d'Italie, 69364 Lyon Cedex 07, France, aguionne@umpa.ens-lyon.fr. Research supported by ANR project ANR-08-BLAN-0311-01.

†Department of Mathematics, UC Berkeley, Berkeley, CA 94720, vfr@math.berkeley.edu. Research supported by NSF grant DMS-0856316.

‡Department of Mathematics, UCLA, Los Angeles, CA 90095 USA, shlyakht@math.ucla.edu. Reserach supported by NSF grants DMS-0555680, DMS-0900776.

We mention also that while this paper was in preparation, Kodiyalam and Sunder have found a different proof that M_k are (in the finite-depth case) isomorphic to interpolated free group factors [KSb].

We conclude the paper with another application of the isomorphism between M_0 and a compression of $\mathfrak{M}$. It allows us to recover the random matrix model used in [GJS08] and can be quite useful in random matrix computations (we illustrate this by describing the joint law of Jones-Wenzl idempotents JW).


2. A SEMI-FINITE TRACIAL ALGEBRA ASSOCIATED TO A PLANAR ALGEBRA.

Let $\mathcal{P}$ be a planar algebra. We denote by $\mathcal{P}_k^\epsilon$, $k = 0, 1, 2, \dots$, $\epsilon = \pm$, the k -th graded component of $\mathcal{P}$.

For fixed k, ϵ and integers a, b, p satisfying $a + b + p = 2k$, let $V_{a,b}^\epsilon(p)$ be a copy of $\mathcal{P}_k^\epsilon$; we think of $V_{a,b}^\epsilon(p)$ as diagrams arranged to have, clockwise from the first string, a strings on the left, p on top, b on the right and so that the top-left corner has shading ϵ . Define the multiplication map


$$\cdot : V_{a,b}^\epsilon(p) \times V_{a',b'}^{\epsilon'}(p') \rightarrow V_{a,b'}^\epsilon(p+p')$$

to be zero unless $b = a'$ and $\epsilon' = (-1)^p \epsilon$ and otherwise by the tangle:



The two choices of shading at the top left of the tangle correspond to the possible values of ϵ .

Define the trace $Tr : V_{a,b}^\epsilon(p) \rightarrow \mathcal{P}_0^\epsilon$ to be zero unless $a = b$ and otherwise by the tangle



(here $\sum TL$ denotes the sum of all Temperley-Lieb diagrams). Finally, consider the inclusions

$$V_{a,b}^\epsilon(p) \rightarrow V_{a+2r,b+2s}^\epsilon(p)$$

given by the tangle

$$(2.3) \quad \delta^{-(r+s)/2} \quad \begin{array}{c} \text{Diagram: A rectangular box containing a tangle. The tangle consists of a large central rectangle with an L-shaped cutout. To the left of the cutout, there are $2r$ loops (represented by three pairs of semi-circles) with a bracket labeled $2r$. To the right of the cutout, there are $2s$ loops (represented by two pairs of semi-circles) with a bracket labeled $2s$. The top of the box has a horizontal bar with three segments, the middle one containing an ellipsis. The left and right sides of the box have vertical bars with three segments each, the middle one containing an ellipsis. There are two asterisks (*) inside the box: one in the upper right area and one in the lower left area. The entire diagram is scaled by $\delta^{-(r+s)/2}$ on the left. \end{array}$$

Let

$$V_{\epsilon_2}^{\epsilon_1}(p) = \bigcup_{\substack{a,b \geq 0 \\ \epsilon_2 = (-1)^{a+b}}} V_{a,b}^{\epsilon_1}(p)$$

and

$$V_{\epsilon_2}^{\epsilon_1} = \bigoplus_{p \geq 0} V_{\epsilon_2}^{\epsilon_1}(p).$$

Finally, we let

$$V_+ = V_+^+ \oplus V_+^-.$$

One easily checks that the inclusions (2.3) are compatible with the multiplication (2.1) and the trace (2.2), thus proving the following:

Lemma 2. (a) Equation (2.1) determines an associative multiplication $\cdot$ on $V_+ = V_+^+ + V_+^-$ so that

$$V_{\epsilon}^{\epsilon'}(p) \cdot V_{\epsilon}^{\epsilon''}(p') \subset V_{\epsilon}^{\epsilon'}(p + p').$$

Furthermore, if $x \in V_+^{\epsilon}(0)$ and $y \in V_+^{\epsilon'}$ and $\epsilon \neq \epsilon'$, then $x \cdot y = 0$.

(b) Equation (2.2) defines a trace on V_+ .

(c) The linear spaces $A_+^{\epsilon} \stackrel{\text{def}}{=} V_+^{\epsilon}(0)$ form subalgebras of V_+ . Moreover, $A_+ = A_+^+ + A_+^-$ is isomorphic to $A_+^+ \oplus A_+^-$ as algebras.

We could also define $V_- = V_-^- \oplus V_-^+$; this is also an algebra with a trace, in the analogous way.


Lemma 3. (a) The tangle

$$(2.4) \quad \begin{array}{c} \text{Diagram: A rectangular box containing a tangle. The tangle consists of a central rectangle. The top of the box has a horizontal bar with three segments, the middle one containing an ellipsis. The left and right sides of the box have vertical bars with three segments each, the middle one containing an ellipsis. There is an asterisk (*) inside the box, centered below the central rectangle. \end{array}$$

defines an injection i from A_+^+ to A_+^- .

(b) The tangle

(2.5)



defines a completely-positive map $E_1 : A_+^- \rightarrow A_+^+$.

(c) Let $\eta : A_+ \rightarrow A_+$ be given by

$$\eta(a \oplus b) = E_1(b) \oplus i(a), \quad a \in A_+^+, b \in A_+^-.$$

Then $\text{Tr}(x\eta(y)) = \text{Tr}(\eta(x)y)$ for all $x, y \in A_+$.

(d) The inclusion of $V_{a,a}^\epsilon(0)$ into $V_{a+2,a+2}^\epsilon(0)$ determined by setting $r = s = 1$ in (2.3) is the same as the inclusion α described in Lemma 2.1 in [PS03]. In particular, the algebra A_+^+ (resp., A_+^-) is exactly (the algebraic inductive limit of $\mathcal{P}_k$'s inside) the type I von Neumann algebra $\mathcal{A}_{-1}^{-1}$ (resp., $\mathcal{A}_0^{-1}$) defined in (2.4.7) in [PS03], and this identification is trace-preserving.

(e) The trace Tr is non-negative definite on A_+ , and moreover there exists a type I semi-finite von Neumann algebra $\mathfrak{A}_+$ with trace Tr containing A_+ as a weakly dense subalgebra in a trace-preserving way. The minimal projections of $\mathfrak{A}_+$ are contained in A_+ .

(f) The minimal central projections of A_+ are labeled by the graph Γ . The trace of a minimal projection of A_+ lying in the central component associated to the vertex v is the value of the Perron-Frobenius eigenvector $\mu(v)$, normalized by $\mu(*) = 1$.

(g) If $\mathfrak{A}_+^\epsilon$ is the closure of A_+^ϵ in $\mathfrak{A}_+$, then $\mathfrak{A}_+ = \mathfrak{A}_+^+ \oplus \mathfrak{A}_+^-$. Moreover, $\mathfrak{A}_+^\epsilon = P_\epsilon \mathfrak{A}_+$, where P_ϵ is the central projection corresponding to all even (if $\epsilon = +$) or odd (if $\epsilon = -$) vertices in the principal graph Γ .


(h) The inclusion $\mathfrak{A}_+^+ \subset \mathfrak{A}_+^-$ is given by the graph Γ .

Proof. Parts (a), (b) and (c), (d) are straightforward. Both (e) and (f) follow from (d); indeed one takes $\mathfrak{A}_+ = \mathcal{A}_{-1}^{-1} \oplus \mathcal{A}_0^{-1}$. The remaining parts (f), (g) and (h) follow from [PS03], Lemma 2.6. $\square$

One could instead work with A_- ; in this case a similar lemma holds, with the exception of replacing Γ by the dual principal graph.

Theorem 4. Let $e_n \in A_+^{(-1)^n}$ denote the projection

(2.6)



Then there are a trace-preserving isomorphisms:

$$\begin{aligned} (e_n V_+ e_n, \cdot, \delta^{-n} Tr) &\cong \begin{cases} (\mathcal{P}, \wedge_n, Tr_n), & n \text{ even}, \\ (\mathcal{P}^{\text{op}}, \wedge_n, Tr_n), & n \text{ odd}; \end{cases} \\ (e_n V_- e_n, \cdot, \delta^{-n} Tr) &\cong \begin{cases} (\mathcal{P}^{\text{op}}, \wedge_n, Tr_n), & n \text{ even}, \\ (\mathcal{P}, \wedge_n, Tr_n), & n \text{ odd}; \end{cases} \end{aligned}$$

where we write $\mathcal{P}^{\text{op}}$ for the dual planar algebra to $\mathcal{P}$ (i.e., one for which the shadings are reversed).

Proof. The isomorphism is given by identifying an element $x \in \mathcal{P}_{2n+p}^\epsilon$ with an element of $V_{n,n}^\epsilon(p)$, and then identifying $V_{n,n}^\epsilon(p)$ with $e_n V_{(-1)^n \epsilon}^\epsilon e_n$. $\square$

3. OPERATOR-VALUED SEMICIRCULAR SYSTEMS.

Let A be a von Neumann algebra and let $\eta : A \rightarrow A$ be a completely-positive map. Then [Shl99, Spe98] there is a unique von Neumann algebra $M \supset A$, a conditional expectation $E : M \rightarrow A$ and an element $X = X^* \in M$ so that: (i) $M = W^*(A, X)$; (ii) $E(a_0 X a_1) = 0$ for any $a_j \in A$ and E satisfies the following recursive property (here $a_1, a_2, \dots \in A$):

$$(3.1) \quad E(a_0 X a_1 \cdots X a_n) = \sum_{k=2}^n a_0 \eta(E(a_1 X a_2 \cdots X a_{k-1})) E(a_k X a_{k+1} \cdots a_{n-1} X a_n).$$


For example,

$$E(a_0 X a_1) = 0, \quad E(a_0 X a_1 X a_2) = a_0 \eta(a_1) a_2.$$

The element X is called the A -valued semicircular element with variance η . Note that η is determined by $\eta(a) = E(X a X)$.

The definition still makes sense if A is a self-adjoint subalgebra of a von Neumann algebra, provided that η extends to a completely-positive map on (some) the von Neumann algebra containing A . The A -valued distribution of X is completely described by the recursive formula (3.1).

It is not hard to see that is equivalent to the following graphical rule of computing $E(a_0 X a_1 \cdots X a_n)$. First draw the product as follows:



Next, consider the following drawing, where $\sum TL$ stands for the sum over all T :

$$(3.2) \quad \begin{array}{c} \boxed{\sum TL} \\ \hline \begin{array}{c} \text{Diagram of } a_0 X a_1 X a_2 \cdots a_{n-1} X a_n \text{ with vertical lines above each } X \end{array} \end{array}$$

Finally, obtain the value of $E(a_0 X a_1 X \cdots a_{n-1} X a_n)$ by recursively performing the following replacements in (3.2):

$$\begin{array}{c} \text{Diagram of } a X b X c \text{ with a curved line connecting the two } X \text{ circles} \end{array} = \boxed{a \eta(b) c}$$

In the case that A is semi-finite with trace Tr and η satisfies

$$Tr(x\eta(y)) = Tr(\eta(x)y), \quad \forall x, y \in \mathcal{L}^1(Tr)$$

the algebra M is also semi-finite with trace $Tr \circ E$.

More generally, a family $\{X_i : i \in I\}$ is called *semicircular over A* if one has the recursive relation

$$E(a_0 X_{i_1} a_1 \cdots X_{i_n} a_n) = \sum_{k=2}^n a_0 \eta_{i_1 i_k} (E(a_1 X_{i_2} a_2 \cdots X_{i_{k-1}})) E(a_k X_{i_{k+1}} a_{k+1} \cdots a_{n-1} X_{i_n} a_n).$$

The joint variance η_{ij} can be viewed as a matrix-valued map $\eta = (\eta_{ij})_{ij} : A \rightarrow A \otimes B(\ell^2(I))$ (where $B(\ell^2(I))$ stands for bounded operators on $\ell^2(I)$), and the positivity requirement is that this map be completely-positive.

We need the following Lemma, which can be found in [Shl99, Shl98]:

Lemma 5. (a) Let X be an A -valued semicircular element, let $a_i, b_i \in A$. Then the elements $\{a_i^* X b_j + b_j^* X a_i\}_{i \leq j}$ form an A -valued semicircular family. (b) Let $X_i : i \in I$ be an A -valued semicircular family. Then X_i are free with amalgamation over A iff $E(X_i a X_j) = 0$ for $i \neq j$ and all $a \in A$. (c) If $A_1 \subset A$ is a subalgebra so that $\eta(A_1) \subset A_1$, then X is also A_1 -semicircular. (d) If $\eta(a) = \tau(a)1$ for all $a \in A$ and a state $\tau : A \rightarrow \mathbb{C}$, then X is free from A in $W^*(A, X, \tau \circ E)$. (e) $\|X\| \leq 2\|\eta(1)\|$.

We now return to the algebra $A_+ = A_+^+ \oplus A_+^-$ that we defined in Lemma 2. Let $\mathfrak{A}_+ = \mathfrak{A}_+^+ \oplus \mathfrak{A}_+^-$ be as in Lemma 3(e). Let $\eta : \mathfrak{A}_+ \rightarrow \mathfrak{A}_+$ be given by

$$\eta(a \oplus b) = E_1(b) \oplus i(a)$$

as in Lemma (2). Then η is a completely-positive map and satisfies $Tr(x\eta(y)) = Tr(\eta(x)y)$ for all $x, y \in \mathcal{L}^1(Tr)$. Let X be an $\mathfrak{A}_+$ -valued semicircular element, and let $\mathfrak{M}_+ = W^*(\mathfrak{A}_+, X)$, $E : \mathfrak{M}_+ \rightarrow \mathfrak{A}_+$ the canonical conditional expectation, and $Tr = Tr \circ E$ a semi-finite trace on $\mathfrak{M}_+$.

Lemma 6. Let $X_n = e_{2n} X e_{2n+1} + e_{2n+1} X e_{2n}$. Let also $f_n = e_{2n} \oplus e_{2n+1} \in A_+$. Then $f_n \uparrow 1$ weakly. Moreover, $X_n = f_n X f_n$, and X_n is an operator-valued semicircular element over $\mathfrak{A}_+$ with variance $\eta_n(x) = f_n \eta(f_n x f_n) f_n$. In particular, since $f_n \mathfrak{A}_+ f_n = f_n A_+ f_n$ and $\eta_n(f_n A_+ f_n) \subset f_n A_+ f_n$, X_n is $f_n A f_n$ -semicircular with variance $\eta|_{f_n A f_n}$.

Consider now the element $c_n \in V_+$ given by the diagram

$$(3.3) \quad c_n = \begin{array}{|c|} \hline \begin{array}{c} \text{[Diagram: A rectangle with a horizontal line near the top. The top-left corner is unshaded, and the rest is shaded. A bracket on the right indicates the height is } 2n+1 \text{.} \end{array} \\ \hline \end{array}$$

(the top-left corner is unshaded).

Lemma 7. (a) $e_{2n} c_n = c_n e_{2n+1}$;
(b) $f_m c_n f_m = c_m$ if $n \geq m$.
(c) If $x_n = c_n + c_n^*$, then $c_n = e_{2n} x_n e_{2n+1}$.

(d) For any N_0 , $V_+ = \text{Alg}(A_+, \{x_n\}_{n \geq N_0})$.


(e) Let $E : V_+ \rightarrow A_+$ be the conditional expectation given by the tangle

(3.4)

Then x_n is A_+ -semicircular with variance $\eta_n(x) = f_n \eta(f_n x f_n) f_n$. In particular, $E(x_n a x_n f_n) = f_n \eta(f_n x f_n) f_n$ for all $a \in A_+$.

Proof. (a), (b), (c) are straightforward verifications.

To prove (d), we clearly have the inclusion $\text{Alg}(A_+, \{x_n\}_{n \geq N_0}) \subset V_+$. Note that c_n and c_n^* belong to $\text{Alg}(A_+, x_n)$. Next, one can obtain any element of V_+ by applying the following tangle to various elements of A_+ :



This tangle, however, can be obtained by composing the multiplication tangle with diagrams of the form

(3.5)

and their adjoints. But the following picture shows that (3.5) (up to the inductive limit defined by (2.3)) belongs to $\text{Alg}(A_+, \{x_n\}_{n \geq N_0})$:

(3.6)

This shows that $\text{Alg}(A_+, \{x_n\}_{n \geq N_0}) = V_+$.

Finally, part (e) follows from the definition of E and (3.2). □

We have thus proved:

Theorem 8. *The map taking A_+ into $\mathfrak{A}_+$ and each x_n to X_n extends to a trace-preserving isomorphism between $(V_+, \cdot, Tr)$ and a dense subalgebra of $\mathfrak{M}_+$. Thus $\mathfrak{M}_+ \cong W^*(\mathfrak{A}_+, X)$, where X is $\mathfrak{A}_+$ -semicircular of variance η .*

Corollary 9. *The trace Tr on V_+ is non-negative definite, and elements of V_+ give rise to bounded operators in the GNS representation on $L^2(V_+, Tr)$.*

Proof. This follows from the fact that $L^2(V_+, Tr) = L^2(\mathfrak{M}_+, Tr)$, and that every element of V_+ acts on $L^2(V_+, Tr)$ in the same way as some finite-degree non-commutative polynomial in X and elements of A_+ . Moreover, $\|X\| \leq 2\|\eta(1)\| < \infty$. $\square$

Corollary 10. *$M_j(\mathcal{P}) \cong e_j \mathfrak{M}_+ e_j$ if j is even and $M_j(\mathcal{P}^{\text{op}}) \cong e_j \mathfrak{M}_+ e_j$ if j is odd. Here $\mathcal{P}^{\text{op}}$ denotes the dual planar algebra and we write $M_j(\mathcal{P})$ for the tower of factors associated to a planar algebra $\mathcal{P}$ in [GJS08].*

Let P_ϵ be the unit of $\mathfrak{A}_+^\epsilon$, $\epsilon = \pm$.

Lemma 11. (a) *Let $c = P_+ X P_-$ and let $c = vb$ be the polar decomposition of c . Then $vv^* = P_+$ and $v^*v \leq P_-$. (b) $P_\pm$ both have central support 1 in $\mathfrak{M}_+$. (c) If the index satisfies $\delta^2 > 1$, $\mathfrak{M}_+$ is a type II_∞ factor.*

Proof. Let $b_n^2 = c_n^* c_n$. Then $b_n \in e_{2n} \mathfrak{M}_+ e_{2n}$ and its law there is the same as the law of $\cup$ in $M_0 \subset M_n$. It follows that if we set $v_n = c_n b_n^{-1/2}$, then $v_n v_n^* = e_{2n}$. Since $e_{2n} \uparrow P_+$, $vv^* = P_+$. On the other hand, since $e_{2n+1} c_n c_n^* e_{2n+1} = c_n c_n^*$, it follows that $v_n^* v_n \leq e_{2n+1}$. Thus $vv^* \leq P_-$. This proves (a). To prove (b), consider a central projection q so that $q \geq P_-$. But then $q \geq v^*v$ and so $q \geq vv^*$ (since q is central). Thus $q \geq P_+$ also, and so $q \geq P_+ + P_- = 1$. Finally, to prove (c), assume that $q \in \mathfrak{M}_+$ is a central projection. Then since $[q, P_\pm] = 0$, we find that $q = P_+ q P_+ + P_- q P_- = q_+ + q_-$, where $q_\epsilon \in P_\epsilon \mathfrak{M}_+ P_\epsilon$. Again, let $f_n = e_{2n} + e_{2n+1}$, $f_n \uparrow 1$. Then $f_n q_\epsilon f_n$ is central in $P_\epsilon \mathfrak{M}_+ P_\epsilon$, which is isomorphic to either $M_{2n+1}(\mathcal{P}^{\text{op}})$ or $M_{2n}(\mathcal{P})$ depending on the parity of ϵ . But these are factors by [GJS08], so $f_n q_\epsilon f_n$ must be multiples of identity. Since $f_n \uparrow 1$, it must be that $q = 0$, $q = P_+$, $q = P_-$ or $q = 1$. But because of (b) only $q = 0$ and $q = 1$ are possible. $\square$

Corollary 12. *If the index satisfies $\delta^2 > 1$, we have the following isomorphisms*

$$M_{2j+1}(\mathcal{P}^{\text{op}}) \cong (M_0)^{\delta^{2j+1}}, \quad M_{2j}(\mathcal{P}) \cong (M_0)^{\delta^{2j}}.$$

Here $\mathcal{P}^{\text{op}}$ is the planar algebra dual to $\mathcal{P}$ and we write $M_j(\mathcal{P})$ for the tower of factors associated to a planar algebra $\mathcal{P}$ in [GJS08].

This is of course because by Corollary 10, $M_1(\mathcal{P}^{\text{op}}) \cong M_0(\mathcal{P})^\delta$ and in general $M_{k+2} \cong M_k^{\delta^2}$.


4. THE ALGEBRA $\mathfrak{M}_+$.

Let Γ be the principal graph of $\mathcal{P}$, and denote by Γ^+ the even and by Γ^- the odd vertices of Γ . For each $v \in \Gamma^+$ choose a minimal projection $q_v \in \mathfrak{A}_+^+$ in the central summand corresponding to v (in the case of $v = *$ we choose $q_* = e_0$). Let μ be the Perron-Frobenius eigenvector for Γ , normalized by $\mu(*) = 1$. Then $Q = \bigoplus_{v \in \Gamma^+} q_v \oplus i(q_v)$ is a projection in $\mathfrak{A}_+$. Moreover, there exists a family of partial isometries $m_i \in \mathfrak{A}_+$ having the form $m_i = w_i \oplus i(w_i)$ and satisfying: $m_i^* m_i = Q$, $\sum m_i m_i^* = 1$.

As in Theorem 8, let X be an $\mathfrak{A}_+$ -semicircular element of variance η .

Lemma 13. *For any $a \in \mathfrak{A}_+^+$, $aX = Xi(a)$.*

Proof. This follows from $aX_n = ac_n$ and the following diagram:



□

Corollary 14. $Q\mathfrak{M}_+Q = W^*(Q\mathfrak{A}_+Q, QXQ)$. Moreover, QXQ is $Q\mathfrak{A}_+Q$ -semicircular with variance $\eta|_{Q\mathfrak{A}_+Q}$.

Proof. $Q\mathfrak{M}_+Q = W^*(Q\mathfrak{A}_+Q, \{m_i^*Xm_j\}) = W^*(Q\mathfrak{A}_+Q, QXQ)$ because of Lemma 13. The rest follows from Lemma 5. □

We now describe the algebra $Q\mathfrak{A}_+Q$. Let us write $Q_+ = QP_+$ and $Q_- = QP_-$. Thus Q_e is the unit of $Q\mathfrak{A}_+Q$. The algebra $Q\mathfrak{A}_+Q$ is abelian. Its minimal (central) projections are the projections q_v , $v \in \Gamma^+$ that we have chosen before. For each edge e of Γ , write $s(e) \in \Gamma^+$ and $t(e) \in \Gamma^-$ to denote the two ends of e . For each $w \in \Gamma^-$ fix a minimal projection $f_w \in Q\mathfrak{A}_+Q$ in the central component associated to w . Next, for each edge e choose a partial isometry $w_e \in Q\mathfrak{A}_+Q$ so that $w_e^*w_{e'} = \delta_{e=e'}f_{t(e)}$ and $\sum_{s(e)=v} w_e w_e^* = i(q_v)$. Let $F = Q_+ + \sum_{w \in \Gamma^-} f_w \leq Q$. Finally, set

$$(4.1) \quad X_e = w_e^* X q_{s(e)} + q_{s(e)} X w_e \in F\mathfrak{M}_+F.$$

Lemma 15. $F\mathfrak{M}_+F = W^*(F\mathfrak{A}_+F, \{X_e : e \in E(\Gamma)\})$, where $E(\Gamma)$ is the set of edges of Γ . Moreover, X_e are free with amalgamation over $F\mathfrak{A}_+F$ and each X_e is $F\mathfrak{A}_+F$ -semicircular with variance η_e given by $\eta_e(a \oplus a') = (E_1(w_e a' w_e^*) \oplus w_e^* i(q_v a q_v) w_e)$, $v = s(e)$.

Proof. Let $F_+ = FQ_+$, $F_- = FQ_-$. Then $w_e^* X q_{s(e)} = F_- X_e F_+$. We now claim that $F\mathfrak{M}_+F$ is generated by $F\mathfrak{A}_+F$ and elements $w_e^* X q_{s(e)}$. Indeed, this follows from the identity $\sum_v \sum_{s(e)=v} w_e w_e^* X q_{s(e)} = Q_- X Q_+$, which allows us to approximate an arbitrary word in $\mathfrak{A}_+$ and X by words in $\mathfrak{A}_+$ and $\{X_e\}_{e \in E(\Gamma)}$. We now apply Lemma 5(a) to $A = Q\mathfrak{A}_+Q$ to conclude that $\{X_e\}_{e \in E(\Gamma)}$ form a semicircular family. Finally, if $e \neq e'$ and $q_v \in F_+ \mathfrak{A}_+ F_+$, $f_w \in F_- \mathfrak{A}_+ F_-$, then

$$\begin{aligned} E(X_e a X_{e'}) &= (w_e^* i(q_{s(e)} q_v q_{s(e')}) w_{e'}) = 0 \\ E(X_e b X_{e'}) &= (q_{s(e)} E_1(w_e f_w w_e^*) q_{s(e')}) = 0, \end{aligned}$$

the last equality because $w_e f_w w_{e'}^* = \delta_{t(e)=w} w_e w_{e'}^*$ and $Tr(w_e w_e^* i(q_v)) = Tr(w_e^* i(q_v) w_e) = Tr(w_e^* \sum w_f w_f^* w_e) = 0$ if $e \neq e'$, so that $E_1(w_f w_e^*) = 0$. Using Lemma 5(b) we conclude that X_e are free with amalgamation over $F\mathfrak{A}_+F$. □

To simplify notation, we shall from now on write $\mathcal{A} = F\mathfrak{A}_+F$, $\mathcal{A}_\epsilon = FQ_\epsilon \mathcal{A}$, $1_\epsilon = FQ_\epsilon$, $\epsilon = \pm$. Note that the algebra $\mathcal{A}$ is abelian, with its minimal central projections labeled by vertices of Γ . We will write p_v for the projection corresponding to $v \in \Gamma$ (thus $p_v = q_v$ if $v \in \Gamma^+$ and $p_w = f_w$ if $w \in \Gamma^-$).

We summarize this as:

Lemma 16. (a) There is a trace-preserving isomorphism $(M_0, Tr_0) \cong (e_0 \mathfrak{N} e_0, Tr)$, where $\mathfrak{N} = F\mathfrak{M}_+F$. (b) $\mathfrak{N}$ is isomorphic to the amalgamated free product $*_{\mathcal{A}} \{\mathfrak{N}_e : e \in E(\Gamma)\}$ where $\mathfrak{N}_e = W^*(\mathcal{A}, X_e)$ and X_e is an $\mathcal{A}$ -valued semicircular element with variance η_e determined by

$\eta_e(p_v \oplus 0) = \lambda_e \delta_{s(e)=v} 0 \oplus p_{t(e)}$, $\eta_e(0 \oplus p_w) = \lambda'_e \delta_{t(e)=w} p_{s(e)} \oplus 0$, where λ_e, λ'_e are some nonzero constant. (c) $\mathfrak{N}_e \cong \bigoplus_{v \in \Gamma \setminus \{s(e), t(e)\}} \mathbb{C} p_v \oplus W^*(p_{s(e)}, p_{t(e)}, X_e)$. (d) If we set $\mathcal{A}_e = \mathbb{C} p_{s(e)} + \mathbb{C} p_{t(e)}$, then X_e is $\mathcal{A}_e$ -semicircular.

Proof. To prove (a), note that $e_0 = q_* \leq F$. Thus $M_0 \cong e_0 \mathfrak{M}_+ e_0 = e_0 F \mathfrak{M}_+ F e_0 = e_0 \mathfrak{N} e_0$ (note that in our normalization $\text{Tr}(e_0) = 1$). Part (b) is nothing by the conclusion of Lemma 15. To see (c), we note that $X_e = (p_{s(e)} + p_{t(e)}) X_e (p_{s(e)} + p_{t(e)})$. Finally, to see (d) we note that η_e takes $\mathcal{A}_e$ to $\mathcal{A}_e$ and apply Lemma 5(c). $\square$

Lemma 17. *Let $e \in E(\Gamma)$ and let $v, w \in \Gamma$ so that $\mu(v) \leq \mu(w)$ and $\{s(e), t(e)\} = \{v, w\}$ (here μ is the Perron-Frobenius eigenvector). Endow the algebra $\mathfrak{N}_e$ with the normalized trace $\tau = \frac{1}{\mu(v) + \mu(w)} \text{Tr}$. Then there exists a trace-preserving isomorphism between $W^*(\mathcal{A}_e, X_e)$ and the algebra $(M_{2 \times 2} \otimes (L^\infty[0, 1])) \oplus \mathbb{C} p$ where the trace on the second algebra is determined by $\tau(p) = (\mu(w) - \mu(v)) / (\mu(w) + \mu(v))$. In particular, the free dimension fdim [BDJ06, Dyk93, Dyk94, Dyk09] of any set of generators of this algebra is given by*

$$\begin{aligned} \text{fdim } W^*(\mathcal{A}_e, X_e) &= 1 - \frac{1}{(\mu(w) + \mu(v))^2} (\mu(w) - \mu(v))^2 \\ &= 1 + \frac{1}{(\mu(w) + \mu(v))^2} (-\mu(v)^2 - \mu(w)^2 + 2\mu(v)\mu(w)) \end{aligned}$$

Proof. Consider the algebra $\mathcal{A}_e$ generated by $p_w \mathbb{C} \oplus p_v \mathbb{C}$ (with unit $p_w + p_v$) and a semicircular element Y which is free from $\mathcal{A}_e$. By Lemma 5(d), Y is $\mathcal{A}_e$ -semicircular with variance τ . Let $X' = p_w Y p_v + p_v Y p_w$. Then X' is $\mathcal{A}_e$ -semicircular and has the same variance (up to a scalar multiple) as X_e .

Thus

$$W^*(\mathcal{A}_e, X_e) \cong W^*(\mathcal{A}_e, X').$$

Let $Z = p_v X' p_w = p_v Y p_w$. Then the distributions of ZZ^* and Z^*Z are free Poisson elements; moreover, ZZ^* has support projection p_v , while Z^*Z has the support projection $q \leq p_w$ of trace $\tau(q) = \tau(p_v)$. Let $Z = V|Z|$ be the polar decomposition of Z . Thus $VV^* = p_v$, $VV^* = q \leq p_w$, so that $W^*(\mathbb{C} p_v \oplus \mathbb{C} p_w, X_e) \cong W^*(p_v, q, V, |Z|, p_w - q)$. Note that the support projection of any element in the algebra generated by $p_v, q, V, |Z|$ is under $p_v + q = 1 - (p_w - q)$, so $W^*(p_v \mathbb{C} \oplus p_w \mathbb{C}, X_e) = W^*(p_v, q, V, |Z|) \oplus \mathbb{C} p$, where $p = p_w - q$ is a projection of trace $\frac{1}{\mu(v) + \mu(w)} (\tau(p_w) - \tau(p_v)) = \frac{1}{\mu(v) + \mu(w)} (\mu(w) - \mu(v))$. On the other hand, since p_v and q are equivalent via V , $W^*(p_v, q, V, |Z|)$ is isomorphic to the algebra of 2×2 matrices over the von Neumann algebra generated by $|Z|$ in the algebra $p_v W^*(p_v \mathbb{C} \oplus p_w \mathbb{C}, Y)$. Since the law of $|Z|$ is quarter-circular and has no atoms, the algebra generated by $|Z|$ (with unit p_v) is isomorphic to $L^\infty[0, 1]$. Thus $W^*(p_v \mathbb{C} \oplus p_w \mathbb{C}, X_e)$ is isomorphic to $(M_{2 \times 2} \otimes (L^\infty[0, 1])) \oplus \mathbb{C} p$ as claimed.

The computation of the “free dimension” fdim can be performed using the formulas in [BDJ06, Dyk93, Dyk94, Dyk09]. Indeed, since the algebra $M_{2 \times 2} \otimes (L^\infty[0, 1])$ is hyperfinite and diffuse, its free dimension is 1. Thus $\text{fdim} [(M_{2 \times 2} \otimes (L^\infty[0, 1])) \oplus \mathbb{C} p] = 1 - \tau(p)^2 = 1 - \frac{1}{(\mu(w) + \mu(v))^2} (\mu(w) - \mu(v))^2$. $\square$

Corollary 18. *$M_0 = e_0 \mathfrak{N} e_0$ where $\mathfrak{N}$ is an amalgamated free product (indexed by edges of Γ) of type I von Neumann algebras with amalgamation over a fixed discrete type I von Neumann subalgebra.*

4.1. The finite-depth case. In the case that Γ is infinite, the projection F is infinite, and the amalgamated free product appearing in Corollary 18 involves an infinity of terms $\mathfrak{N}_e$, each semifinite. Unfortunately, we have been unable to find existing results in the literature to handle this case, although it is natural to conjecture that the resulting factor is then the infinite amplification of $L(\mathbb{F}_\infty)$. However, in the case that Γ is finite, the projection F is finite and so $\mathfrak{N}$ is a finite factor. Moreover, each term $\mathfrak{N}_e$ is also finite and is of type I. In this case one can apply the results of and formulas in [BDJ06, Dyk93, Dyk94, Dyk09].

Lemma 19. *There is a trace-preserving isomorphism $(\mathfrak{N}, \frac{1}{Tr(F)}Tr) \cong (L(\mathbb{F}(s)), \tau)$ where*

$$(4.2) \quad s = 1 + \frac{1}{Tr(F)^2} \left(- \sum_{v \in \Gamma} \mu(v)^2 + 2 \sum_{e \in E(\Gamma)} \mu(t(e))\mu(s(e)) \right),$$

$$Tr(F) = \sum_{v \in \Gamma} \mu(v).$$

Proof. The algebra $\mathfrak{N}$ is an amalgamated free product over all edges e of Γ of the algebras $\mathfrak{N}_e$ over the subalgebra $\mathcal{A}$. Let $T = \sum_{v \in \Gamma} \mu(v) = Tr(F)$.

One has $\text{fdim}(\mathcal{A}) = 1 - T^{-2} \sum_{v \in \Gamma} \mu(v)^2$. Furthermore,

$$\begin{aligned} \text{fdim}(\mathfrak{N}_e) &= \text{fdim} \left(\bigoplus_{v \in \Gamma \setminus \{t(e), s(e)\}} \mathbb{C}p_v \oplus W^*(\mathcal{A}_e, X_e) \right) \\ &= 1 - \left(\sum_{v \in \Gamma \setminus \{t(e), s(e)\}} T^{-2} \mu(v)^2 \right) - T^{-2} \left(\mu(s(e)) - \mu(t(e)) \right)^2 \\ &= 1 - \left(\sum_{v \in \Gamma} T^{-2} \mu(v)^2 \right) + T^{-2} \left(\mu(s(e))^2 + \mu(t(e))^2 \right) \\ &\quad - T^{-2} \left(\mu(t(e))^2 + \mu(s(e))^2 - 2\mu(s(e))\mu(t(e)) \right) \\ &= \text{fdim}(\mathcal{A}) + 2T^{-2} \mu(s(e))\mu(t(e)). \end{aligned}$$

Let e_0 an edge starting from $*$. By Lemma 11(c), $\mathfrak{N} = F\mathfrak{M}_+F$ is a factor.

Then by [BDJ06, Dyk93, Dyk94, Dyk09], since $\mathfrak{N}$ is a factor, it is isomorphic to $L(\mathbb{F}(s))$ where

$$\begin{aligned} s &= \text{fdim} \mathfrak{N}_{e_0} + \sum_{e \in E(\Gamma) \setminus \{e_0\}} (\text{fdim} \mathfrak{N}_e - \text{fdim}(\mathcal{A})) \\ &= \text{fdim} \mathcal{A} + 2T^{-2} \mu(s(e_0))\mu(t(e_0)) + 2T^{-2} \sum_{e \in E(\Gamma) \setminus \{e_0\}} \mu(s(e))\mu(t(e)) \\ &= 1 + \frac{1}{T^2} \left(- \sum_{v \in \Gamma} \mu(v)^2 + 2 \sum_{e \in E(\Gamma)} \mu(t(e))\mu(s(e)) \right). \end{aligned}$$

□

Theorem 20. *If $\mathcal{P}$ is finite depth with global index I and $\delta > 1$, then (a) $M_0 \cong L(\mathbb{F}(1 + 2(\delta - 1)I))$, (b) $M_k \cong L(\mathbb{F}(1 + 2\delta^{-2k}(\delta - 1)I))$.*

Proof. By Corollary 18, $M_0 \cong e_0 \mathfrak{N} e_0$, where $e_0 \in \mathfrak{N}$ is a projection of trace $1/\text{Tr}(F) = 1/T$ in the notation of Lemma 19. Moreover, $\mathfrak{N} = L(\mathbb{F}(s))$ with s given by (4.2).

Recall that the eigenvector condition implies that $\sum_{e:s(e)=v} \mu(v)\mu(t(e)) = \delta \sum \mu(v)^2$. From this we find that $\sum_{e \in E(\Gamma)} \mu(s(e))\mu(t(e)) = \delta \sum_{v \in \Gamma^+} \mu(v)^2$. Let I be the global index [Ocn88, EK98], $I = \sum_{v \in \Gamma^+} \mu(v)^2$. Thus $\sum_{e \in E(\Gamma)} \mu(s(e))\mu(t(e)) = \delta I$.

The eigenvector condition implies that $\sum_{v \in \Gamma} \mu(v)^2 = 2 \sum_{v \in \Gamma^+} \mu(v)^2 = 2I$. Hence

$$s = 1 + \frac{1}{T^2} \left(\sum_{v \in \Gamma} \mu(v)^2 + 2 \sum_{e \in E(\Gamma)} \mu(s(e))\mu(t(e)) \right) = 1 + \frac{1}{T^2} (-2I + 2\delta I) = 1 + 2T^{-2}(\delta - 1)I.$$

The compression formula [Răd94, Dyk94, VDN92] then implies that $M_0 \cong L(\mathbb{F}(r))$ with $r = 1 + T^2(s - 1) = 1 + 2(\delta - 1)I$. This proves (a). For k even, part (b) follows from the compression formula and Corollary 12. In the odd case, applying our arguments to $\mathcal{P}^{\text{op}}$ instead of $\mathcal{P}$ and using the compression formula and Corollary 12, we get that $M_{2k+1}(\mathcal{P}^{\text{op}}) \cong L(\mathbb{F}(1 + 2\delta^{-2k}(\delta - 1)I^{\text{op}}))$ where I^{op} is the global index of $\mathcal{P}^{\text{op}}$. But $I^{\text{op}} = I$ (since the principal graphs of $\mathcal{P}$ and $\mathcal{P}^{\text{op}}$ share a bipartite half) and so (b) holds. $\square$

4.2. Some examples.

4.2.1. Finite-dimensional Kac algebras. We first consider the case of the planar algebra of an n -dimensional Kac algebra as considered in [KSa]. In this case, the principal graph Γ has the form

$$* \text{ --- } \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \end{array} \right\} (n-1) \text{ vertices.}$$

The Perron-Frobenius eigenvector is equal to 1 on all even vertices and to $\sqrt{n}$ on the single odd vertex. The associated eigenvalue is $\delta = \sqrt{n}$. The global index is given by $I = n \cdot 1 = n$. In this case, Theorem 20 gives:

$$M_0 \cong L(\mathbb{F}(1 + 2n\sqrt{n} - 2n)), \quad M_1 \cong L(\mathbb{F}(2\sqrt{n} - 1)),$$

in accordance with the results of [KSa].

4.2.2. The inclusion $M \subset M \otimes M_{n \times n}$. Another example is the graph planar algebra associated to the graph $* \equiv \bullet$ (n edges). The associated inclusion of II_1 factors is of the form $M \subset M \otimes M_{n \times n}$. In this case, $\delta = n$, the eigenvector is equal to 1 at all vertices, and the global index is 1. Thus $M_0 \cong L(\mathbb{F}(2n - 1))$.

5. $\mathfrak{N}_+$ AND RANDOM BLOCK MATRICES.

We point out a connection between our description of $M_0 = e_0 \mathfrak{N}_+ e_0 = e_0 W^*(\mathcal{A}, \{X_e : e \in E(\Gamma)\}) e_0$ and certain random block matrices considered in §3.1 of [GJS08]. To avoid the clash of notation, we will write $\bar{X}_e$ for what was denoted X_e in that paper:

Proposition 21. *The limit joint distribution of the random matrices $\{d_v : v \in \Gamma\} \cup \{\bar{X}_e + \bar{X}_{e^{\text{op}}}^* : e \in E(\Gamma)\}$ as defined in [GJS08, §3.1] is the same as that of $\{p_v : v \in \Gamma\} \cup \{X_e : e \in E(\Gamma)\} \subset W^*(\mathcal{A}, \{X_e : e \in E(\Gamma)\}) = \mathfrak{N}_+$.*

Proof. The variables X_e constructed in the present paper form an $\mathcal{A}$ -valued semicircular family, and so do the variables $\bar{X}_e + \bar{X}_e^*$ constructed in [GJS08] (see the proof of Proposition 2). $\square$

In fact, there is a clear similarity between our construction of X_e from the variable X given by equation (4.1) and the construction of Y_e in §3.3 of [GJS08].

We now recall the realization of $(\mathcal{P}, \wedge_0, Tr_0)$ constructed in [GJS08]. First, a planar algebra $\mathcal{P}$ is realized as the subalgebra of $\mathcal{P}^\Gamma$ associated to its principal graph. Let us call this inclusion ι . For any $w \in \mathcal{P}$ one can then write $\iota(w) = \sum_\rho w_\rho \rho$, where the summation takes place over all loops on Γ starting at an even vertex, identified with elements of $\mathcal{P}^\Gamma$. If we then write $X_\rho = X_{e_1} \cdots X_{e_n}$ in the case that $\rho = e_1 \cdots e_n$, $e_j \in E(\Gamma)$, one of the main results of [GJS08] states that the map

$$\pi : w \mapsto \sum_\rho w_\rho \bar{X}_\rho$$

is a $*$ -homomorphism from $(\mathcal{P}, \wedge_0, Tr_0)$ to the von Neumann algebra generated by $\{p_v : v \in \Gamma\}$ and $\{X_e : e \in E(\Gamma)\}$, and that π satisfies

$$E_{W^*(p_v : v \in \Gamma^+)}(\pi(w)) = \sum_{v \in \Gamma^+} Tr_0(w) p_v,$$

where E denotes the Tr -preserving conditional expectation.

We now make the observation that the map

$$\pi_0 : w \mapsto e_0 \pi(w) e_0 = \sum_{\rho \text{ starting at } *} w_\rho \bar{X}_\rho$$

is once again a $*$ -homomorphism from $(\mathcal{P}, \wedge_0, Tr_0)$, which this time satisfies $Tr(\pi_0(w)) = Tr_0(w)$. But modulo the identifications given by Proposition 21, this is exactly the isomorphism between $(\mathcal{P}, \wedge_0, Tr_0)$ and $e_0 \mathfrak{N}_+ e_0$.

5.1. A short alternative proof of Theorem 20 for M_0 . We note that one can *deduce* Proposition 21 from the fact that π_0 intertwines Tr_0 and Tr , thus giving an alternative (and very short) proof to the fact that $M_0 \cong e_0 \mathfrak{N}_+ e_0$, where $\mathfrak{N}_+$ is *defined* to be $\mathfrak{N}_+ = W^*(\mathcal{A}, X_e : e \in E(\Gamma))$. One can then proceed as in §4.1, thus obtaining a shorter way of identifying the isomorphism class of M_0 . We have taken the longer route in this paper for the purpose of introducing and exploring the extra structure of the II_∞ factor $\mathfrak{M}_+$.

5.2. Computations of moments of Jones-Wenzl projections JW_n . We now consider case of the TL planar algebra, whose principal graph is A_K , $K = 2, 3, \dots, \infty$.

Up to normalization, each simple path (i.e., a path which is a geodesic between $*$ and the farthest point from $*$ that it reaches) of length $2n$ corresponds to a Jones-Wenzl projection in $\mathcal{P}_n$. Thus in the A_n case we get a single Jones-Wenzl projection JW_k for each $k = 1, 2, \dots, K-1$. If we denote by e_j the j -th edge in the graph A_K , $1 \leq j \leq K-1$ (in our numbering, e_1 starts at $*$), then JW_k corresponds to the path $e_1 \dots e_k e_k^o \dots e_1^o$.

Thus the joint law of the Jones-Wenzl projections is the same as that of $JW_1 = \bar{X}_{e_1} \bar{X}_{e_1}^*$, $JW_2 = \bar{X}_{e_2} \bar{X}_{e_1} \bar{X}_{e_1}^* \bar{X}_{e_2}^*$, $JW_3 = \bar{X}_{e_3} \bar{X}_{e_2} \bar{X}_{e_1} \bar{X}_{e_1}^* \bar{X}_{e_2}^* \bar{X}_{e_3}^*$ and so on.

In particular, let us use this to compute the law ν_n of JW_n inductively. We have that (writing $c_n = P_+ X_{e_n} P_-$ and $Q_n = c_1 \cdots c_n = Q_{n-1} c_n$):

$$(5.1) \quad \int t^k d\nu_n(t) = \text{Tr}_0((JW_n)^k) = \text{Tr}((Q_n Q_n^*)^k) = \text{Tr}((c_n c_n^* Q_{n-1}^* Q_{n-1})^k).$$

Here Tr_0 is the finite trace on $\mathcal{P}, \Lambda_0$ and Tr is the semi-finite trace on $\mathfrak{N}_+$. Note that both $c_n c_n^*$ and Q_{n-1} belong to $p_n \mathfrak{N}_{+p_n}$, where we write p_n for the projection at which e_n starts. Let $\mu_n = \text{Tr}(p_n)$. Then $\mu_n^{-1} \text{Tr}$ is a normalized trace on $p_n \mathfrak{N}_{+p_n}$. Moreover, we see from the random matrix model (in which c_k is modeled by an independent block matrix of size $\mu_k N \times \mu_{k+1} N$, $N \rightarrow \infty$) that if u is a Haar unitary free from c_n and Q_{n-1} with respect to $\mu_n^{-1} \text{Tr}$, then (5.1) does not change if we replace c_n by $u c_n$. It follows that $c_n c_n^*$ and $Q_{n-1}^* Q_{n-1}$ are free in $(p_n \mathfrak{N}_{+p_n}, \mu_n^{-1} \text{Tr})$, and so

$$\nu_n = (\text{law of } c_n c_n^* \text{ with respect to } \mu_n^{-1} \text{Tr}) \boxtimes (\text{law of } Q_{n-1}^* Q_{n-1} \text{ with respect to } \mu_n^{-1} \text{Tr}).$$

We first compute the law of $Q_{n-1}^* Q_{n-1}$. To this end, we note that

$$\mu_n^{-1} \text{Tr}((Q_{n-1}^* Q_{n-1})^k) = \mu_n^{-1} \text{Tr}((Q_{n-1} Q_{n-1}^*)^k) = \mu_n^{-1} \int t^k d\nu_{n-1}(t).$$

Thus the law of $Q_{n-1}^* Q_{n-1}$ is given by $\mu_n^{-1} \nu_{n-1} + (1 - \mu_n^{-1}) \delta_0$.

To compute the law of $c_n c_n^*$ we note that it is (using the random matrix model) the Marčenko-Pastur distribution associated to a $\mu_n N \times \mu_{n+1} N$ rectangular matrix, $N \rightarrow \infty$. In other words, it is the Free Poisson law π_λ with parameter $\lambda = \mu_{n+1}/\mu_n$ and thus has S -transform

$$S_n(z) = \frac{1}{z + \mu_{n+1}/\mu_n}.$$

Thus (noting that $\mu_1 = 1$ in our normalization)

$$\nu_n = \pi_{\mu_{n+1}/\mu_n} \boxtimes (\mu_n^{-1} \nu_{n-1} + (1 - \mu_n^{-1}) \delta_0), \quad \nu_1 = \pi_{\mu_2}.$$

Let us now denote by Ξ_n the S -transform of ν_n . Then the S -transform of $\mu_n^{-1} \nu_{n-1} + (1 - \mu_n^{-1}) \delta_0$ is given by

$$\Xi_{n-1}(\mu_n z) \frac{1+z}{\mu_n^{-1} + z} = \Xi_{n-1}(\mu_n z) \frac{\mu_n + \mu_n z}{1 + \mu_n z}$$

Thus

$$\Xi_n(z) = \Xi_{n-1}(\mu_n z) \frac{\mu_n(\mu_n + \mu_n z)}{(1 + \mu_n z)(\mu_{n+1} + \mu_n z)}, \quad \Xi_1(z) = \frac{1}{z + \mu_2}.$$

REFERENCES

- [BDJ06] N. Brown, K. Dykema, and K. Jung, *Free entropy dimension in amalgamated free products*, preprint math.OA/0609080, 2006.
- [Dyk93] K. Dykema, *Free products of hyperfinite von Neumann algebras and free dimension*, Duke Math J. **69** (1993), 97–119.
- [Dyk94] ———, *Interpolated free group factors*, Pacific J. Math. **163** (1994), 123–135.
- [Dyk09] ———, *A description of amalgamated free products of finite von Neumann algebras over finite dimensional subalgebras*, Preprint, arXiv.org/0911.2052, 2009.
- [EK98] D. Evans and Y. Kawahigashi, *Quantum symmetries on operator algebras*, Oxford Univ. Press, New York, 1998.
- [GJS08] A. Guionnet, V. Jones, and D. Shlyakhtenko, *Random matrices, free probability, planar algebras and subfactors*, To appear in Proceedings NCG, arXiv.org/0712.2904, 2008.

- [KSa] V. Kodyalam and V. S. Sunder, *Guionnet-Jones-Shlyakhtenko subfactors associated to finite-dimensional Kac algebras*, Preprint, arXiv.org:0901.3180.
- [KSb] ———, *On the Guionnet-Jones-Shlyakhtenko construction for graphs*, Preprint arXiv.org:0911.2047.
- [Ocn88] A. Ocneanu, *Quantized group, string algebras and Galois theory for algebras*, Operator algebras and applications, London Math. Soc. Lect. Notes Series, vol. 136, Cambridge University Press, 1988, pp. 119–172.
- [PS03] S. Popa and D. Shlyakhtenko, *Universal properties of $L(\mathbf{F}_\infty)$ in subfactor theory*, Acta Math. **191** (2003), no. 2, 225–257. MR MR2051399 (2005b:46140)
- [Răd94] F. Rădulescu, *Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index*, Invent. math. **115** (1994), 347–389.
- [Shl98] D. Shlyakhtenko, *Some applications of freeness with amalgamation*, J. reine angew. Math. **500** (1998), 191–212.
- [Shl99] ———, *A-valued semicircular systems*, J. Func. Anal **166** (1999), 1–47.
- [Spe98] R. Speicher, *Combinatorial theory of the free product with amalgamation and operator-valued free probability theory*, Mem. Amer. Math. Soc. **132** (1998), x+88.
- [VDN92] D.-V. Voiculescu, K. Dykema, and A. Nica, *Free random variables*, CRM monograph series, vol. 1, American Mathematical Society, 1992.